

Literaturliste - Subchondrale Knochenzysten

von Rechenberg B, Guenther H, McIlwraith CW, et al: Fibrous tissue of subchondral cystic lesions in horses produce local mediators and neutral metalloproteinases and cause bone resorption in vitro. Vet Surg 2000; 29:420-429.

McIlwraith CW: Subchondral cystic lesions (osteochondrosis) in the horse. Compend Contin Educ Pract Vet 1982;4:s282-294.

Baxter GM: Subchondral cystic lesions in horses, in McIlwraith CW, Trotter GW (eds): Joint disease in the horse. Philadelphia,PA, Saunders, 1996, pp 384–397.

Jackson M, Fricker C, Kümmerle J und Fürst A: Die Behandlung von subchondralen zystoiden Defekten beim Pferd mit Benzopyron: Eine retrospective Analyse (2008). Wien. Tierärztl. Mschr. 95 (2008), 158 – 165.

Wallis TW, Goodrich LR, McIlwraith CW, et al. Arthroscopic injection of corticosteroids into the fibrous tissue of subchondral cystic lesions of the medial femoral condyle in horses: A retrospective study of 52 cases (2001-2006). Equine Vet J 2008;40:461-467



Jackson WA, Stick JA, Arnoczky SP, et al: The effect of compacted cancellous bone grafting on the healing of subchondral bone defects of the medial femoral condyle in horses. Vet Surg 2000;29:8-16

Kraus KH, Kirker-Head C: Mesenchymal stem cells and bone regeneration. Vet Surg 2006;35:232-242

Ortved KF, Nixon AJ, Mohammed HO, et al: Treatment of subchondral cystic lesions of the medial femoral condyle of mature horses with growth factor enhanced chondrocyte grafts: A retrospective study of 49 cases. Equine Vet J 2012;44:606-613

Fürst AE, Kaegi B, von Rechenberg B, et al: Die Behandlung von 5 Pferden mit subchondralen zystoiden Defekten im Fesselbein. Pferdeheilkunde 1997;13:147-161.

Jackson MA, Olerth S, Kummer M, et al: Use of a parathyroid hormone peptide (PTH1-34)-enriched fibrin hydrogel for the treatment of subchondral cystic lesions in 15 horses. Pferdeheilkunde 2012;28:1-6.

Santschi EM, Williams JM, Morgan JW, et al: Preliminary Investigation of the Treatment of Equine Femoral Condylar Subchondral Cystic Lesions With a Transcondylar Screw. Vet Surg 2014;1-8.

SEEUFER MEDIENPRODUKTION OG

Große Pfarrgasse 4/8, 1020 Wien, Tel.: +43 699 8148 5455, E-Mail: redaktion@news4vets.at
Firmenbuchnummer: FN 368685i, Firmenbuchgericht: Handelsgericht Wien, UID-ATU66744847
Raiffeisenbank Weiden am See, IBAN: AT383310600000018119, BIC: RLBBAT2E106



Smith MA, Walmsley JP, Phillips TJ, et al: Effect of age at presentation on outcome following arthroscopic debridement of subchondral cystic lesions of the medial femoral condyle: 85 horses (1993-2003). Equine Vet J 2005;37:175-180.

Kuemmerle J, Auer JA, Rademacher N, Lischer C et al: Short Incomplete Sagittal Fractures of the Proximal Phalanx in Ten Horses Not Used for Racing. Vet Surg 2008; 193-200.

Kirker-Head C.A., Boudrieau R.J., Kraus K.H. Use of bone morphogenetic proteins for augmentation of bone regeneration. JAVMA 231:1039-1055, 2007.

McKay W.F. and Peckham S.M. A comprehensive clinical review of recombinant human bone morphogenetic protein-2 (INFUSE® Bone Graft). Int Orthopaedics (SICOT) 31:729-734, 2007.

SEEUFER MEDIENPRODUKTION OG

Große Pfarrgasse 4/8, 1020 Wien, Tel.: +43 699 8148 5455, E-Mail: redaktion@news4vets.at
Firmenbuchnummer: FN 368685i, Firmenbuchgericht: Handelsgericht Wien, UID-ATU66744847
Raiffeisenbank Weiden am See, IBAN: AT383310600000018119, BIC: RLBBAT2E106

